

## Solubility of Low Volatile Hydrocarbons in Supercritical Ethane

R. Eustaquio-Rincon

*Programa de Ingeniería Molecular, Instituto Mexicano del Petróleo, México D. F., Mexico*

A.F. Paz-Menendez

*Programa de Ingeniería Molecular, Area de Investigación en Termofísica, Instituto Mexicano del Petróleo, México D. F., Mexico*

A. Trejo<sup>C,S</sup>

*Competencia de Ciencias Básicas, Instituto Mexicano del Petróleo, México D. F., Mexico*

Supercritical fluid processes are being used in a lot of applications, including hydrocarbon systems, such as for fractionation of aromatic heavy oil [1], extraction of petroleum hydrocarbons from soil [2], etc. This is due to the special properties of supercritical solvents. Carbon dioxide is an option for extraction of heavy hydrocarbons [3] since CO<sub>2</sub> has many advantages: it is a green solvent, it has a low critical temperature, it is non-flammable, non-toxic, economic, *etc.* In addition, the solubility of heavy hydrocarbons could be improved with the use of cosolvents or entrainers [4]. However, it is important to evaluate other solvents, like light hydrocarbons, in order to establish the best option for extracting heavy hydrocarbons.

In this work we report the solubility of n-octacosane in ethane and squalane in ethane both at 308 and 323 K, in the pressure range 10 to 20 MPa. The experimental studies were carried out in a flow apparatus. The extracted solute was determined gravimetrically and the solubility data was determined to  $\pm 0.0002$  mole fraction. The solubility of the hydrocarbons increases as pressure increases but the functionality with pressure shows a special behavior. The solubility data were correlated with the Chrastil equation through density values of ethane.

From a comparison of the solubility of n-octacosane in ethane at 323 K with solubility data reported in the literature with CO<sub>2</sub> at the same temperature [5] it was possible to establish that the solubility of the hydrocarbon in ethane is larger two orders of magnitude than that of CO<sub>2</sub>. This result clearly shows that the like dissolves the like.

[1] S.J. Park, C.J. Kim, and B.S. Rhee. Ind. Eng. Chem. Res. 4897-7900, 39(2000).

[2] S. Liang and D.C. Tilotta. Anal. Chem. 616-622, 70(1998).

[3] R. Eustaquio-Rincón and A. Trejo. Fluid Phase Equilib. 231-239, 185(2001).

[4] R. Eustaquio-Rincón and A. Trejo. Submitted to the Fifteenth Symposium on Thermophysical Properties.

[5] M.A. McHugh, A.J. Seckner, and T.J. Yogan. Ind. Eng. Chem. Fundamen. 493-499, 23(1984).